

Reply to “Proof of the Gauge Independence of the Conformal Anomaly of Bosonic String in the Sense of Kraemmer and Rebhan”

Ulrike Kraemmer and Anton Rebhan

*Institut für Theoretische Physik, Technische Universität Wien,
Wiedner Hauptstr. 8-10, A-1040 Vienna, Austria*

February 1, 2008

Abstract

In two recent preprints (hep-th/9710131 and 9710132), Abe and Nakanishi have claimed that the proof of the gauge independence of the conformal anomaly of the bosonic string as given by us in 1988 was wrong. A similar allegation has been made concerning our proof of the gauge independence of the sum of the ghost number and Lagrange multiplier anomalies in non-conformal gauges. In this short note we refute their criticism by explaining the simple logic of our proofs and emphasizing the points that have been missed by Abe and Nakanishi.

The standard approach to covariant quantization of string theory chooses the conformal gauge, which in the critical dimension permits to eliminate the world-sheet metric as a dynamical variable[1]. The gauge independence of the conformal anomaly, which determines the critical dimension, has been investigated first in Ref. [2], but only for algebraic gauges. In Ref. [3], the world-sheet anomalies were explicitly calculated for the harmonic gauge, which appeared to be of particular interest because of a vanishing ghost-number anomaly. The conformal anomaly on the other hand remained unchanged as expected. In Ref. [4] the present authors have confirmed these results, but have shown that the eliminated ghost-number anomaly has just been shifted to an analogous number-current anomaly for the now dynamical world-sheet metric; the sum of the ghost-number and the so-called Lagrange-multiplier anomalies turned out to be gauge independent. The gauge independence of the conformal anomaly was also verified in Refs. [5, 6, 7], which considered the background-covariant harmonic gauge, and in Refs. [8, 4], where non-background-covariant de Donder gauges were used.

In Ref. [9] we have reviewed and extended the explicit calculations of the various world-sheet anomalies and have given a simple proof of the observed gauge independences based on BRS symmetry. In a more general framework, the gauge independence of anomalies was subsequently confirmed in Refs. [10, 11].

Some years later, a seemingly contradictory result was published by Abe and Nakanishi[12], who claimed that in non-conformal gauges the conformal anomaly was undetermined due to ambiguities in the definition of the energy-momentum tensor.¹ Although later Ref. [13] has pointed out that their ambiguity disappears if one refrains from restricting to a flat background prematurely, in Refs. [14, 15] Abe and Nakanishi have recently questioned the previous works that showed the gauge independence of world-sheet anomalies. In particular they claimed that the explicit proof in Ref. [9] was based on false assumptions and therefore wrong. (Not trusting the more general arguments of Refs. [10, 11], they presented an alternative proof which is however so restricted that they cannot even “compare two gauges which have different Feynman rules”.)

In Ref. [9] we have considered diffeomorphism and Weyl gauge conditions

¹The explicit result of Ref. [12] is in fact erroneous. As Abe and Nakanishi had to admit recently [15], they had overlooked infrared divergences which restrict their alleged ambiguity to a sign ambiguity.

of the form

$$F_i^{mn}[\hat{g}]h_{mn} = 0, \quad \Omega^{mn}[\hat{g}]h_{mn} = 0, \quad i, m, n = 0, 1 \quad (1)$$

where $h_{mn} = g_{mn} - \hat{g}_{mn}$ with $\hat{g}$ being a classical background field. Imposing these gauge conditions by Lagrange multipliers b^i and b , respectively, one finds that h_{mn} and (b^i, b) fields have only mixed propagators. As a consequence, the one-loop effective action is exact and the gauge-fixed action can be linearized with respect to the *quantum* metric field h_{mn} . Correspondingly, the BRS symmetry can be linearized, which moreover implies abelianization.

As is well known, while anomalies arise only in the context of renormalization, they are contained already in the regularized effective action Γ , which is a functional of the background metric field $\hat{g}_{mn}$. Local contributions to the latter can be changed by renormalization, but non-local ones can not. Using this freedom to restore any diffeomorphism invariance that a particular regularization scheme may have violated [16], the effective action is proportional to $\int d^2x \sqrt{-\hat{g}} \hat{R} \hat{\square}^{-1} \hat{R}$, and its prefactor determines the conformal anomaly. Since however everything is fixed by the non-local contributions, one only needs to consider the regularized effective action in order to establish the gauge independence of the conformal anomaly.

An infinitesimal variation δF of the gauge condition F can be shown to give

$$\delta\Gamma[\hat{g}] = \left\langle Q_{\text{BRS}}(\bar{c}^i \delta F_i^{mn} h_{mn}) \right\rangle \quad (2)$$

where only the linearized BRS charge is needed as explained above. For the linearized BRS transformations there cannot appear any anomalies, hence (2) is zero and the conformal anomaly is gauge-fixing independent.

In Ref. [14], Abe and Nakanishi remark firstly that this argument cannot be correct, for it would imply that the anomaly itself was zero, not only its gauge variations, if δF was replaced by F . They do not provide further explanations, so one can only guess that they have in mind that the gauge sector of the gauge fixed action is BRS exact and that therefore one could delete the δ 's in (2). But the effective action is not the expectation value of the classical action in the sense of (2). Nor could one integrate (2) starting from $F \equiv 0$, because there the effective action would no longer be well-defined.

More concretely, Abe and Nakanishi then point out that variations of the background field and BRS transformations do not commute. This is true, but contrary to what they claim, our proof does *not* depend on this.

In order to determine the proportionality factor in

$$\Gamma[\hat{g}] \propto \int d^2x \sqrt{-\hat{g}} \hat{R} \hat{\square}^{-1} \hat{R} \quad (3)$$

it is sufficient to extract its bilinear terms in an expansion of $\hat{g}_{mn} = \eta_{mn} + \hat{h}_{mn}$. No more and no less is done in the explicit calculations in Refs. [3, 4, 9].

The very same strategy is followed in our treatment [9] of number-current anomalies. These anomalies are contained in an effective action augmented to include external sources coupled to the various number currents. Their first variation with respect to these sources give functionals $\langle \mathcal{J}_i \rangle [\hat{g}]$ whose nonlocal contributions cannot be changed by local counterterms. It is therefore sufficient to study their gauge dependences. Here it turns out that there are gauge dependences, but the sum of the gauge-parameter variations of the ghost number and the Lagrange multiplier currents can again be written as vacuum insertion of a linearized-BRS-exact operator, which explains the findings of the explicit calculations [3, 4, 9].

On the other hand, what Abe and Nakanishi have been doing was to look at correlators of energy-momentum tensors. They appear to be unconcerned with any renormalization issues and therefore are in no position to assess the ambiguities they believe to have found. Their alternative “proof” of gauge independence of the conformal anomaly is therefore rather a collection of technical observations, which even as such appear to be extremely limited.

There are certainly alternative approaches to studying world-sheet anomalies, we have merely chosen one according to our preferences. For instance, in Ref. [17] the conformal anomaly has more recently been determined for a non-linear de Donder gauge,² by considering the anomaly of the (full) BRS symmetry that occurs if one insists on conformal invariance. As is well known, BRS symmetry can be restored by local counterterms at the expense of conformal invariance. This standard procedure has also been criticized by Abe and Nakanishi in Ref. [15], who apparently find it impossible to accept that BRS symmetry could ever be violated. Again, they seem to disregard the connection between the existence of anomalies and issues of renormalization.

To summarize, we find the various criticisms put forward in Refs. [14, 15] completely baseless.

²It is called harmonic gauge in Ref. [17], but in the terminology of our Ref. [9] we have reserved this term for the background-covariant version.

References

- [1] A.M. Polyakov, Phys. Lett. **B103** (1981) 207
- [2] S. Hwang and R. Marnelius, Nucl. Phys. **B272** (1986) 389
- [3] D.W. Düsedau, Phys. Lett. **B188** (1987) 51
- [4] A. Rebhan and U. Kraemmer, Phys. Lett. **B196** (1987) 477
- [5] J.I. Latorre, Nucl. Phys. **B297** (1988) 171
- [6] D.Z. Freedman, J.I. Latorre and K. Pilch, Nucl. Phys. **B306** (1988) 77
- [7] V. Del Duca, L. Magnea, P. van Nieuwenhuizen, Int. J. Mod. Phys. **A3** (1988) 1081
- [8] L. Baulieu and A. Bilal, Phys. Lett. **B192** (1987) 339
- [9] U. Kraemmer and A. Rebhan, Nucl. Phys. **B315** (1989) 717
- [10] W. Kummer, Phys. Lett. **B231** (1989) 53;
O. Piguet, D. Schwarz and M. Schweda, Phys. Lett. **B245** (1990) 393
- [11] W. Kummer, “Gauge independence of anomalies”, in *Fields and Particles*, eds. H. Mitter and W. Schweiger (Springer-Verlag, 1990) pp. 231–253; Acta Phys. Polonica **22** (1991) 37
- [12] M. Abe and N. Nakanishi, Mod. Phys. Lett. **A7** (1992) 1799
- [13] W. Mödritsch, Mod. Phys. Lett. **A9** (1994) 241 ([hep-th/9308074](#))
- [14] M. Abe and N. Nakanishi, preprint RIMS-1161 ([hep-th/9710131](#))
- [15] M. Abe and N. Nakanishi, preprint RIMS-1164 ([hep-th/9710132](#))
- [16] A. Rebhan, U. Kraemmer and R. Knienider, Phys. Rev. **D39** (1989) 3625
- [17] T. Takahashi, Prog. Theor. Phys. **95** (1996) 995 ([hep-th/9512015](#))